

Monopoles in lattice Electroweak theory

B.L.G. Bakker^a, A.I.Veselov^b, M.A. Zubkov^b

^a *Department of Physics and Astronomy, Vrije Universiteit, Amsterdam, The Netherlands*

^b *ITEP, B.Chermushkinskaya 25, Moscow, 117259, Russia*

Abstract

There exist several types of monopole - like topological defects in Electroweak theory. We investigate properties of these objects using lattice numerical methods. The intimate connection between them and the dynamics of the theory is established. We find that the density of Nambu monopoles cannot be predicted by the choice of the initial parameters of Electroweak theory and should be considered as the new external parameter of the theory. We also investigate the difference between the versions of Electroweak theory with the gauge groups $SU(2) \otimes U(1)$ and $SU(2) \otimes U(1)/Z_2$. We do not detect any difference at $\alpha \sim \frac{1}{128}$. However, such a difference appears in the unphysical region of large coupling constant $\alpha > 0.1$.

In both cases we use the following lattice variables: 1. The gauge field $\mathcal{U} = (U, \theta)$, where $U \in SU(2)$, $e^{i\theta} \in U(1)$ realized as link variables. 2. A scalar doublet Φ_α , $\alpha = 1, 2$. The potential for the scalar field is considered in its simplest form in the London limit, i.e., in the limit of infinite bare Higgs mass. From the very beginning we fix the unitary gauge $\Phi_1 = \sqrt{\gamma}$, $\Phi_2 = 0$. For the case of the $SU(2) \times U(1)/Z_2$ symmetric model we chose the action of the form (A) $S_g = \beta \sum_{\text{plaquettes}} ((1 - \frac{1}{2} \text{Tr } U_p \cos \theta_p) + \frac{1}{2} (1 - \cos 2\theta_p)) + \gamma \sum_{xy} (1 - \text{Re}(U_{xy}^{11} e^{i\theta_{xy}}))$, where the plaquette variables are defined as $U_p = U_{xy} U_{yz} U_{wz}^* U_{xw}^*$, and $\theta_p = \theta_{xy} + \theta_{yz} - \theta_{wz} - \theta_{xw}$ for the plaquette composed of the vertices x, y, z, w . For the case of the conventional $SU(2) \times U(1)$ symmetric model we use the action (B) $S_g = \beta \sum_{\text{plaquettes}} ((1 - \frac{1}{2} \text{Tr } U_p) + 3(1 - \cos \theta_p)) + \gamma \sum_{xy} (1 - \text{Re}(U_{xy}^{11} e^{i\theta_{xy}}))$. The following variables are considered as creating a Z boson and a W boson, respectively: $Z_{xy} = Z_x^\mu = \sin [\text{Arg } U_{xy}^{11} + \theta_{xy}]$, $W_{xy} = W_x^\mu = U_{xy}^{12} e^{-i\theta_{xy}}$. Here, μ represents the direction (xy) . In the unitary gauge there is also a $U(1)$ lattice gauge field, which is defined as $A_{xy} = A_x^\mu = [-\text{Arg } U_{xy}^{11} + \theta_{xy}] \bmod 2\pi$. The phase diagrams of the two models under consideration are presented in figure 1. The dashed vertical line represents the phase transition in the $SU(2) \otimes U(1)$ -symmetric model. This is the confinement-deconfinement phase transition corresponding to the $U(1)$ constituents of the model. The same transition for the $SU(2) \otimes U(1)/Z_2$ -symmetric model is represented by the solid vertical line. The dashed horizontal line corresponds to the transition between the broken and symmetric phases of model A. The continuous horizontal line represents the same transition in model B. Interestingly, in the $SU(2) \otimes U(1)/Z_2$ model both transition lines meet, forming a triple point. Real physics is commonly believed to be achieved within the phases of the two models situated in the right upper corner of Fig. 1. The double-dotted-dashed vertical line on the right-hand side of the diagram represents the line, where the renormalized α is constant and equal to $1/128$. All simulations were performed on lattices of sizes 8^4 and 16^4 . Several points were checked using a lattice 24^4 . In general we found no significant difference between the mentioned lattice sizes.




Figure 1: The phase diagrams of the models in the (β, γ) -plane.

We perform the calculation of renormalized fine structure constant α_R using the potential for infinitely heavy external fermions. We consider Wilson loops for the right-handed external leptons: $\mathcal{W}_{\text{lept}}^R(l) = \langle \text{Re} \Pi_{(xy) \in l} e^{2i\theta_{xy}} \rangle$. Here l denotes a closed contour on the lattice. We consider the following quantity constructed from the rectangular Wilson loop of size $r \times t$: $\mathcal{V}(r) = \lim_{t \rightarrow \infty} \log \frac{\mathcal{W}(r \times t)}{\mathcal{W}(r \times (t+1))}$. Due to exchange by virtual photons at large enough distances we expect the appearance of the Coulomb interaction $\mathcal{V}(r) = -\frac{\alpha_R}{r} + \text{const.}$

The worldlines of the quantum Nambu monopoles could be extracted from the field configurations as follows: $j_Z = \delta\Sigma = \frac{1}{2\pi} * d([dZ'] \bmod 2\pi)$ (The notations of differential forms on the lattice are used here.) The monopole density is defined as $\rho = \left\langle \frac{\sum_{\text{links}} |j_{\text{link}}|}{4L^4} \right\rangle$, where L is the lattice size. In order to investigate the condensation of monopoles we use the percolation probability $\Pi(A)$. It is the probability that two infinitely distant points are connected by a monopole cluster. We show Nambu monopole density and percolation probability as a function of γ along the line of constant renormalized $\alpha_R = 1/128$. It is clear that the percolation probability is the order parameter of the transition from the symmetric to the broken phase¹. We also measure the magnetic energy (both $SU(2)$ and $U(1)$), which is carried by Nambu monopoles. The behavior of ΔS_p shows that a quantum Nambu monopole may indeed be considered as a physical object.

In order to evaluate the mass of the Z -boson we use the zero - momentum correlator: $\sum_{\vec{x}, \vec{y}} \langle \sum_{\mu} Z_x^{\mu} Z_y^{\mu} \rangle \sim e^{-M_Z |x_0 - y_0|} + e^{-M_Z (L - |x_0 - y_0|)}$. Here the summation $\sum_{\vec{x}, \vec{y}}$ is over the three “space” components of the four - vectors x and y while x_0, y_0 denote their “time” components. L is the lattice length in the “time” direction. The physical scale is given in our lattice theory by the value of the Z -boson mass $M_Z^{\text{phys}} \sim 91$ GeV. Therefore the lattice spacing is evaluated to be $a \sim [91 \text{ GeV}]^{-1} M_Z$, where M_Z is the Z boson mass in lattice units. The real continuum physics should be approached along the the line of constant $\alpha_R = \frac{1}{128}$, i.e. along the line of constant physics. We investigated the dependence of the ultraviolet cutoff $\Lambda = a^{-1} = (91 \text{ GeV})/M_Z$ on γ along the line of constant physics. It occurs that Λ is increasing slowly along this line with decreasing γ and achieves the value 430 ± 40 GeV at the transition point between the physical Higgs phase and the symmetric

¹The hypothesis that Nambu monopole condensation accompanies the Electroweak transition was first suggested in [3]. Z -vortices condensation in 3D lattice model of high temperature Electroweak theory was investigated numerically in [2].




Figure 2: Nambu monopole density and percolation probability as a function of γ along the line of constant $1/\alpha_R = 128$.

phase. According to our results this value does not depend on the lattice size. This means that the largest achievable value of the ultraviolet cutoff is equal to 430 ± 40 GeV if the potential for the Higgs field is considered in the London limit.

Our lattice study also demonstrates another peculiar feature of Electroweak theory. If we are moving along the line of constant $\alpha = 1/128$, then the Nambu-monopole density decreases with increasing γ (for $\gamma > 1$). Its behavior is approximated with a nice accuracy by the simple formula: $\rho \sim e^{2.08-4.6\gamma}$. This means that the density of Nambu monopoles in the continuum theory cannot be predicted by the choice of the usual parameters of the Electroweak theory and should be considered as a new external parameter of the theory.

We found that the two definitions of the theory (with the gauge groups $SU(2) \otimes U(1)/Z_2$ and $SU(2) \otimes U(1)$, respectively) do not lead to different predictions at the values of α around $1/128$. However, the corresponding models behave differently at unphysically large values of $\alpha > 0.1$. The main difference is in the behavior of the so-called hypercharge monopoles.

This work was partly supported by RFBR grants 08-02-00661, and 07-02-00237, RFBR-DFG grant 06-02-04010, by Grant for leading scientific schools 679.2008.2, by Federal Program of the Russian Ministry of Industry, Science and Technology No 40.052.1.1.1112.

References

- [1] Y. Nambu, Nucl.Phys. B **130**, 505 (1977);
Ana Achucarro and Tanmay Vachaspati, Phys. Rept. **327**, 347 (2000); Phys. Rept. **327**, 427 (2000).
- [2] M.N. Chernodub, F.V. Gubarev, E.M. Ilgenfritz, and A. Schiller, Phys. Lett. B **434**, 83 (1998);
M.N. Chernodub, F.V. Gubarev, E.M. Ilgenfritz, and A. Schiller, Phys. Lett. B **443**, 244 (1998).
- [3] M.N. Chernodub, JETP Lett. **66**, 605 (1997)
- [4] B.L.G. Bakker, A.I. Veselov, and M.A. Zubkov, Phys. Lett. B **583**, 379 (2004);
- [5] B.L.G. Bakker, A.I. Veselov, and M.A. Zubkov, Yad. Fiz. **68**, 1045 (2005).
- [6] B.L.G. Bakker, A.I. Veselov, and M.A. Zubkov, Phys. Lett. B **620**, 156 (2005)

- [7] B.L.G. Bakker, A.I. Veselov, and M.A. Zubkov, Phys. Lett. B **642**, 147 (2006).
- [8] M.I. Polikarpov, U.J. Wiese, and M.A. Zubkov, Phys. Lett. B **309**, 133 (1993).
- [9] B.L.G. Bakker, A.I. Veselov, and M.A. Zubkov, Phys. Lett. B **471**, 214 (1999).
- [10] M.A. Zubkov, Phys. Lett. B **649**, 91 (2007).
- [11] W.Langguth, I.Montvay, P.Weisz Nucl.Phys.B277:11,1986.
- [12] W. Langguth, I. Montvay (DESY) Z.Phys.C36:725,1987
- [13] Anna Hasenfratz, Thomas Neuhaus, Nucl.Phys.B297:205,1988